

Complex network analysis of wind tunnel experiments on the passive scalar dispersion in a turbulent boundary layer

Original

Complex network analysis of wind tunnel experiments on the passive scalar dispersion in a turbulent boundary layer / Iacobello, Giovanni; Ridolfi, Luca; Marro, Massimo; Salizzoni, Pietro; Scarsoglio, Stefania. - ELETTRONICO. - (2019), pp. 65-66. (Intervento presentato al convegno iTi CONFERENCE ON TURBULENCE VIII tenutosi a Bertinoro (Italia) nel 04/09/2018 - 07/09/2018).

Availability:

This version is available at: 11583/2713216 since: 2018-09-17T15:42:40Z

Publisher:

Örlü, Talamelli, Oberlack, Peinke

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Complex network analysis of wind tunnel experiments on the passive scalar dispersion in a turbulent boundary layer

G. Iacobello ¹, L. Ridolfi ², M. Marro ³, P. Salizzoni ³, and S. Scarsoglio ¹

¹ Department Mechanical and Aerospace Engineering, Politecnico di Torino, Turin, Italy

² Department Environmental, Land and Infrastructure Engineering, Politecnico di Torino, Turin, Italy

³ Laboratory of Fluid Mechanics and Acoustics, Ecole Centrale de Lyon, Ecully, France

key-words: Wind tunnel experiments, Complex networks

Abstract:

The characterization of pollutant transport is crucial for the analysis and modeling of the atmospheric boundary layer [1]. In this work, experimental data of passive scalar plumes in a turbulent boundary layer are investigated. The experiments are performed in a wind tunnel where the passive scalar is injected through an L-shaped tube located at $z/\delta = 0.24$, with δ the boundary layer thickness. Specifically, two source configurations are analyzed for two different tube diameters, $D = 3$ mm and $D = 6$ mm. The streamwise, u , and wall-normal, w , velocity components, and the concentration, c , are measured at different distances from the source and at various wall-normal and spanwise locations. The corresponding time-series are recorded for $T = 180$ s with a sampling frequency of 1000 Hz. A sketch of the experimental setup and representative time-series at fixed spatial location is reported in Fig. 1a.

By exploiting the recent advances of complex networks theory, the velocity and concentration time-series are mapped into networks, where each time-sample corresponds to a node. Different algorithms to construct a network from a time-series are explored. One of the simplest and most adopted approaches is the *visibility algorithm* [2, 3]. In a visibility-based network a link exists if the straight line connecting two data-samples lies above the other in-between data. Accordingly, a visibility network is built for each velocity and concentration time-series at different measurement locations (e.g., see Fig. 1b). The resulting network topology inherits the temporal features of the mapped time-series, revealing non-trivial information of the underlying transport process. Due to their versatility, complex networks offer a powerful and synthetic tool for the analysis of turbulent transport and mixing.




Figure 1: (a) Sketch of the wind tunnel set-up and representative velocity and concentration time-series; (b) Networks built from the time-series shown in panel (a).

References

- [1] Nironi, C., Salizzoni, P., Marro, M., Mejean, P., Grosjean, N., and Soulhac, L., (2015), Dispersion of a passive scalar fluctuating plume in a turbulent boundary layer. Part I: Velocity and concentration measurements, *Boundary-layer meteorology*, vol. 156(3), pp. 415-446.
- [2] Gao, Z. K., Small, M., and Kurths, J., (2017), Complex network analysis of time series, *Europhysics Letters*, vol. 116(5), pp. 50001.
- [3] Iacobello, G., Scarsoglio, S., and Ridolfi, L., (2018), Visibility graph analysis of wall turbulence time-series, *Physics Letters A*, vol. 382(1), pp. 1-11.